

Physiotherapy interventions evidence table – Managing psychological factors

The following table provides a summary of level I or II evidence (according to the NHMRC evidence hierarchy) for physiotherapy-relevant interventions in RA published between January 2012 and June 2015. Interpreting the evidence can be complex. RAP-eL users should consider the following:

- There are no current studies investigating the effects of managing psychological factors in early versus late rheumatoid arthritis.
- Further research is needed into the optimal content, format, mode of delivery (face to face, internet, phone), duration and frequency of programmes aimed at improving management of psychological factors.
- It is important to note that the interventions studied are done so in isolation, so the evidence refers to the effect of the single intervention, and not the effect of a multimodal intervention.

Physiotherapy-related intervention(s)	Sources of evidence (see key below)	Results	Making sense of the evidence								
Mindfulness training	<table border="1"> <tr> <td>RCT</td> <td>SR</td> <td>MA</td> <td>CSR</td> </tr> <tr> <td>✓</td> <td></td> <td></td> <td></td> </tr> </table> <p>Fogarty FA et al. Ann Rheum Dis 2015;74:472–474 [PubMed link]</p>	RCT	SR	MA	CSR	✓				<p>Mindfulness based training can:</p> <ul style="list-style-type: none"> • Improve psychological wellbeing in people with RA. • Reduce disease activity as measured by DAS28, morning stiffness and pain scores. 	<ul style="list-style-type: none"> - Mindfulness-based meditation may be a suitable co-intervention for people with RA, delivered via trained facilitators either 1-1 or in a small group format. - For more information and practice tools on mindfulness, refer to painHEALTH.
	RCT	SR	MA	CSR							
	✓										
<table border="1"> <tr> <td>RCT</td> <td>SR</td> <td>MA</td> <td>CSR</td> </tr> <tr> <td>✓</td> <td></td> <td></td> <td></td> </tr> </table> <p>Zangi HA et al. Ann Rheum Dis. 2012;71:911–17. [PubMed link]</p>	RCT	SR	MA	CSR	✓						
RCT	SR	MA	CSR								
✓											
<table border="1"> <tr> <td>RCT</td> <td>SR</td> <td>MA</td> <td>CSR</td> </tr> <tr> <td>✓</td> <td></td> <td></td> <td></td> </tr> </table> <p>Pradhan E et al. Arthritis Care Res 2007;57:1134–42.</p>	RCT	SR	MA	CSR	✓						
RCT	SR	MA	CSR								
✓											

	[PubMed link]					
Managing depression in patients with rheumatoid arthritis (RA)	RCT	SR	MA	CSR	<ul style="list-style-type: none"> • Cognitive behavioural therapy significantly reduced depression in 2 studies. • There is some evidence (Wang, 2008; Neuberger, 2007) to show reduced depression with physical activity-based treatment when compared with control interventions (no exercise/usual care Neuberger (2007); an RA education and stretching exercise group of comparable length to the exercise/tai chi intervention in Wang, 2008). 	<ul style="list-style-type: none"> - Cognitive behavioural therapy and physical activity can both be considered for managing depression in patients with RA. - Further research is required regarding the dose and content of such programmes.
				✓		
	Cramp et al 2013 Non-pharmacological interventions for fatigue in Rheumatoid Arthritis [link]					
	RCT	SR	MA	CSR		
✓						
Wang (2008) Med Sport Sci; 52: 218-229. [link]						
RCT	SR	MA	CSR			
✓						
Neuberger (2007) Arthritis Care & Res; 57(6): 943-952. [link]						

Key To Evidence Sources:

- Randomised Controlled Trial (RCT)
- Systematic Review (SR)
- Meta-Analysis (MA)
- Cochrane Systematic Review (CSR)

List of Table Abbreviations:

- ADL's – Activities of Daily Living
- DAS28 – Disease activity score calculator for Rheumatoid arthritis [\[click here for link to PDF\]](#)
- DASH – “Disabilities of the Arm Shoulder and Hand” outcome measure

RAP-eL

HEP – Home Exercise Programme

HRQ – Health Risk Questionnaire

JP – Joint Protection

LBP – Lower Back Pain

OA – Osteoarthritis

OT – Occupational Therapy

QOL – Quality Of Life

RA – Rheumatoid Arthritis

RCT – Randomised Controlled Trial

TENS – Transcutaneous Electrical Nerve Stimulation

US - Ultrasound

1st MTPJ – 1st Metatarsophalangeal Joint